



Genetically Modified Foods: Pros and Cons for Human Health

Fatemeh Karami^{1*}, Peyman Mahasti Shotorbani²

¹ Applied Biophotonic Research Center, Science and Research Branch, Islamic Azad University, Tehran, Iran

² Department of Food Quality Control and Hygiene, Science and Research Branch, Islamic Azad University, Tehran, Iran

Abstract

Review Article

Received Date:

Oct.15.2017

Revised Date:

Nov.10.2017

Accepted Date:

Nov.28.2017

✉ Correspondent**Author:**

Dr. Fatemeh Karami
Applied Biophotonic
Research Center, Science
and Research Branch,
Islamic Azad University,
Tehran, Iran
fateme.karami@gmail.com

Genetically modified foods (GM foods) have revolutionized the agricultural industry towards more proficient farming in every geographical region of the world. The major tasks of this GM foods technology could be summarized in four points including increase in food products, more convenient food processing, disease prevention and treatment and avoiding of using pesticides through the generation of pest-resistant crops. Although GM food technology had great advantages for human health, there are some concerns regarding biodiversity induced by modified plants, which can indirectly affect human being, as well. Further studies are warranted to define precisely the status of this technology in ecosystem in which human is living.

Keywords: Genetically modified foods, Agriculture, Human health

Introduction

Genetically modified foods which are also known as GM foods or bioengineered foods are foods which are generated from organisms that their genome have been undergone genetic manipulations and modifications to induce a novel trait in order to improve the quality or quantity of product. Although GM foods sometimes refer to the products from organisms along with genetic engineering, genetically modified organisms (GMOs) specifically stands for genetic engineering of animals to produce edible products with novel or improved feature. History of GM foods was officially begun since 1994 by introduction of delayed-ripening

tomato (known as Flavr Savr) which got received its approve on 1992 from food and drug administration (FDA)¹. However, the exact and real time of beginning of GM foods production may be returned to 10,500 to 10,100 BC when the farmers caused selective breeding through artificial selection and making crossing over between plants with genes encoding desired traits and characteristics². By the discovery of DNA structures and genes in later years, the possibility of directed engineering of various types of plants had been provided to reach the desired trait.

GM food industry had been worked on many different agricultural products with major aims of the followings:



1-1 Increasing food production

Enhancement of food productions was one of the most important reasons that led to the introduction of GM food technology. The bitter fact behind this reason is the poor agricultural economy in developing and under developing countries due to have no access and enough money to buy pesticides to protect their products against pest and microbial contaminations³. Moreover, with increasing in world's population within the next half century especially in developing countries, the demand for more productive and efficient food and agricultural industry will be more felt as well in spite of decrease in water and early requirements supplies⁴. Therefore, developing an strong agricultural industry not only can fulfill the future nutrition needs and prevent malnutrition, but also can provide more and desired and various types of unique products using less and inexpensive substrate and less plant treatment.

1-2 Food processing

Plants ripening rate is one of the most important factors in plant delivery and export and can indirectly affect the economy of a country. In this regard, the first delayed ripening plant which has been introduced by Calgene Company in 1994 was tomato (Flavr Savr). It was made through insertion of polygalacturonase inhibitor oligonucleotide into the genome of tomato to delay the ripening process. It is of note that polygalacturonase which is also known as pectin depolymerase degrades galacturonic acid within the pectin network and plays critical role in smoothing and sweetening of fruits and plants as well as tomato⁵. In reverse, accelerated ripening of some dairy products such as Cheddar cheese through over expression of enzymes involved in ripening process has shown a remarkable improvement in dairy industry and economy⁶.

The same scenario has been introduced to double the growth rate of salmon fish through over expression of growth hormone which has get its approval from FDA as the primary GMO (Aqu Advantagea Salmon)⁷.

1-3 Using of GM foods as disease prophylaxis

One of the promising and interesting features of GM foods is genetic engineering in order to enhance the nutritional value of diet in usual dietary sharing schedule. This is very important strategy in developing countries wherein in most of the times almost of the families eat an especial type of crop as well as rice or maize with specific and limited nutritional values which will definitely be associate with increased risk of malnutrition³. This aim can be achieved through two general ways: the first one is known as biofortification with the main purpose of prevention of malnutrition and increasing the nutritional value of present foods.

There are two major strategy in food fortification including direct addition of desired nutrients in to the fertilizer and selective breeding of plant with or without directed mutated plants using selectable markers^{8,9}.

In the first strategy, successful fortification would consider the distribution of nutrient as well as minerals using the appropriate method and soil contents. As an instance, in the districts with Zn deficient soils, using of Zinc (Zn) fertilizers (such as ZnSO₄) which are highly mobile within the soil resulted in production of Zn rich cereals and grains with special trend in more Zn content in some specific plant genotypes¹⁰. The same scenario is held for selenium and iodine as highly mobile mineral nutrients which have been introduced in selenium and iodine poor soils of Finland and New Zealand^{11,12}. The other example is using of Nitrogen Phosphorous and Potash Potassium (NPK) fertilizer in regions with poor soil of



those nutrients which can effectively spread to the plant and enhance its growth (Figure 1)¹³. NPK fertilizer also can be used to increase the spread of other nutrients with low mobility such as iron (Fe) through acidification of soil in the case of NH_4 as an alternative to using of iron chelators as well as FeSO_4 sprays¹³. The major drawback of this strategy is the need for regular and expensive application of fertilizers, which may be also dangerous for ecosystem. The other disadvantage is that it could be helpful for specific types and strains of grain and therefore cannot prevent of all types of malnutrition.




Figure 1: Comparison the growth of the same plants harvested with NPK (left) and without NPK (right).

The second strategy is based on the selective breeding of plants with desired genetic composition to have more potent and rich products in next generations. This strategy is itself can be applied using two main policies which the first one is breeding with plants with dominant genotypes encoding higher amounts of nutrients and minerals. In this regard, a cooperative project was began by Consultative Group on International Agricultural Research (CGIAR) to screen the plants (maize, wheat, cassava, rice and bean) with genotypes encoding high amounts of Fe and Zn¹⁴⁻¹⁶.

The second policy is directed genetic manipulation of plant genome to overexpress the desired protein or other micro or

macromolecules. Target proteins are almost trace element and metal binding proteins and transporter proteins that are abundant in the parts of plant which are not usually edible^{17,18}. Genetic engineering of interest plant can be performed using different approaches including direct transfer of desired gene or direct manipulation of target plant genome using novel genetic engineering technologies as well as CRISPR-Cas9 system and indirect methods usually through making directed infection using bacterial cells containing plasmid to transfer and replicate the desired gene or genes in plant¹⁹⁻²¹.

One of the famous trace elements binding proteins is leghemoglobin, which has pivotal role in concentrating the iron reserve in highly absorbable hem form. Overexpression of this protein in legumes and induced its transcription in other plants as well as cereals can have dramatic effect on iron bioavailability in diets enriched with iron absorption inhibitors. However, direction of targeted protein expression to the edible parts of plants seems to be unsuccessful in its initial attempts and requires further investigations²². The other introduced strategies include insertion of phytase gene to break down the phytate as an inhibitory factor of trace element absorption²³, induced expression of vitamins with enhancing effects on metals and trace elements absorption as well as vitamin C or β -carotene²⁴ and overexpression of proteins containing amino acids which increase the bioavailability of trace elements as well as cysteine^{25,26}.

1-4 Implication of GM foods in pharmacology and disease treatment

Another GM foods technology has relied on production of edible vaccine to not only increase the effect of it but also alleviate the possible side effects of injectable vaccines. Plant based vaccine technology can overcome



on the production and selling circle including the specific storage condition and costs which usually limit appropriate service in some regions and on biosafety concerns, as well²⁷. Interestingly, this strategy has been applied on other organism as well as fishes, which has provided effective, cost benefit and easy to use vaccination²⁸. In fact, every part of plant including fruit, stem, root and etc., can be used to enhance the immune system of human host against various diseases especially infectious diseases by providing required proteins, vitamins and other nutrients. This aim can be fulfilled by cloning the interest antigen encoding gene in to the desired part of plant which will be edible and marketed. One of the amazing advantages of this strategy is that the introduced antigen will be enclosed in within the cellular wall of plant against enzymatic degradation and therefore leads to higher level of immunization²⁹. The other interesting point of plant based vaccination is that those vaccines will be introduced in the human body when the plant reaches the small intestine wherein antigens can be specifically absorbed and directed through special mechanisms through activation of both B and T cells responses^{30,31}. However, some studies have demonstrated that edible vaccines may cause oral tolerance to introduced antigen due to presentation of antigens to T cells by immature dendritic cells. In production of plant derived vaccine against hepatitis B virus (HBV) this problem has been overcome through using of carrier proteins as well as core protein (HBc Ag) as adjuvant or initiating the vaccination process by intramuscular administration to enhance the immune system excitation^{32,33}. Among human vaccine, producing plant derived malaria vaccine had been evaluated in many studies for more than one decade. Although, the frequency and mortality rate of malaria has been remarkably decreased, demand for cost effective and easy using

vaccine with minimum rate of side effects and enough immune response in oral administration is still strongly felt³⁴.

The other amazing examples of plant derived vaccine were potato and corn which were genetically engineered to express enterotoxigenic *E. coli* (ETEC) toxin called as LT-B which is standing for B subunit of it. The advantage of this type of vaccine was described to excite the immune response against glycolipids of plant cell membrane bound with LT-B without inducing diarrhea and thereby immunize the patient against *E. coli* contamination in a safe, easy and relatively cheap manner³⁵. The same procedure has been performed on norovirus, influenza and hepatitis B and even human immunodeficiency virus (HIV) through expression of immunogenic capsid protein within the virus-like particles (VLP) which is not infective in transgenic tobacco leaves, potato tubers, lettuce, tobacco, tomato, carrot, rice, maize and Arabidopsis³⁶⁻⁴². There are other plant-based vaccines which are some of them are in the final phases of clinical trials including H5 pandemic influenza, seasonal influenza, Ebola Virus, Hepatitis C virus (HCV) and human papillomavirus (HPV)⁴³⁻⁴⁷. Anthrax which is an infection caused by *Bacillus anthracis* and frequently involves skin and intestine has been shown to be effectively preventive in mice and rabbit by genetic engineering of *Nicotiana benthamiana* as vaccine⁴⁸.

Another medicine-based application of GM foods is improvement in enzyme replacement therapy (ERT) which had been successful in getting food and drug administration (FDA) approval. One of the best examples is ERT of Gaucher's disease in which the patients have hereditary deficiency of the enzyme glucocerebrosidase actively involved in metabolism of glucocerebroside. In this regard, the primary FDA approval has been devoted to the novel Plant molecular farming (PMF)



approach marketed by Protalix and used to induce the expression of the same enzyme, recombinant taliglucerase alpha, in transgenic carrot cells^{49,50}. The second enzyme deficiency disease is Fabry disease, which is as a result of defective metabolism of sphingolipids (α -Galactosidase-A deficiency) and their accumulation in nervous system and other parts of body. PRX-102 is a novel enzyme with the same structure and function of α -Galactosidase-An enzyme, which has been expressed and marketed by Protalix with significant stability compared to commercial enzyme through genetic engineering of tobacco, and *Nicotiana benthamiana* Leaves are passing through the phase I and II of clinical trials^{45,51}. Plants based medicines have seized great attentions to be used in cancer immunotherapy in other ways than dietary administration. Plant aided immunotherapy has been fulfilled through using almost of plant viruses as well as potexvirus PVX as carrier of antigen targeting tumor cells within the structure of drug conjugates to efficiently excite immune response against tumor cells^{52,53}.

1-5 Resistance to insect and infectious agents

Almost all of plants are susceptible to be infected or damaged with different types of insects and plant specific infections such as fungi, nematodes, viruses and bacteria. Those infections not only threaten the agricultural economy of a country but also sometimes may have dangerous side effects on human health. In this regard, making foods with higher stability and tolerance against contamination including over expression or inducing resistance of herbicide target proteins and enhancing the expression of proteins responsible for degradation of substantial herbicide factors, had dramatic improvement in cultivar plants especially crops. The initial trials have been performed by Calgene company in 1987 with

introducing tobacco with tolerance to herbicides containing enzymes as well as glyphosate, bromoxynil and sulfonylurea^{54,55}. The technology of herbicide resistant (HR) crops has been developed since 1996 to efficiently control the weeds. Weeds usually compete with plants in using water, nutrients and especially space of growth and thereby constrain the production of their interest targets. Recently, Iquebal *et al.* have primarily studied gene expression profile of chickpea to identify the important genes associated with tolerance to herbicide Imazethapyr. Chickpea is belonged to legume crops which is a comparable source of protein relative to animal products⁵⁶. Shoba *et al.* were successful in making directed mutagenesis in rice called as HTM-N22 which was tolerant to Imazethapyr, as well⁵⁷. The most recent approach in developing pest resistant plants is inhibitory RNA (iRNA) mediated suppression of pests as well as Aphids (Aphididae)⁵⁸. The mechanism of action of all the iRNAs is suppression of target RNA translation through forming RNA induced silencing complex (RISC). This is a natural defense against viral infection in most of the Eukaryotes as well as plants which can be directed against any other pest or parasite by designing the nucleotide sequence of miRNA complementary to the major target gene encoding protein of parasite⁵⁹.

There are many reports conferring using of many different approaches in producing plants resistant to insects through cloning of gene encoding of toxin derived from different bacterial cells. As an instance, transgenic rice encoding *Bacillus thuringiensis* (Bt) toxin has shown to be successful in protecting them against attack of spiders with no side effect on spider's population⁶⁰. Another example is insect resistant maize called as TC1507 which has been transfected with modified Cry1F and PAT genes and has been approved to be cultivated in US since 2001⁶¹.



1- What is behind of GM foods consumption
GM foods and GMO had major improvement in agricultural technologies including overproduction of plants in less time, no further need to use pesticides with carcinogenic effects, possibility of growing plants in every geographical region with every soil composition, help to save fossil fuels and less CO₂ emission. The mentioned major agricultural and environmental advantages of GM food technology have indirectly improved the human health as well as edible vaccines, which can remarkably influence the disease incidence and their prevention especially in developing countries. Despite of those benefits and advantages there are some other concerns regarding taking GM foods, which have been noted in different studies. Those concerns can be discussed in two ways including the effect on first environment and ecosystem and the second with more importance on human health.

One of the main studied aspects of ecosystem affected by GM foods is changing in food web of some animals as well as arthropod especially in long term assay^{62,63}. As an instance, some studies have revealed that Bt maize has raised the frequency of other non target organisms which can indirectly affect the arthropod mortality rate⁶⁴. The other aspect which can be influenced by GM foods is gene flow through breeding between GM and non-GM plants which is itself can be affected by different biotechnological and environmental factors⁶⁵. In this case, gene flow not only results in unwanted contamination of non-GM plants but also the generated hybrid plants hybridized usually are less fit compared to wild ones⁶⁶. Although it seems that natural selection is the final determinant of detrimental or favorite allele's frequency within an agricultural land, generation of transient invasive or abnormal plants may have crucial impacts on regional economy. The chemistry of soil including PH, phosphorous, potassium and organic matters

has been shown to be changed in area cultivated with GM plants versus non-GM plants⁶⁷.

Regarding the effects of GM foods with the aim of prevention or treatment of the diseases on human health, we should keep in our mind that every drug with FDA approve has its own adverse effect and uncontrolled consumption of even natural foods and plants has definitely side effects which may threaten health status of peoples. However, despite of many investigations performed on animal studies, there is no report indicating no histopathologic or clinical side effects⁶⁸. However, every genetic modification has a chance to be associated with unintended helpful or harmful mutations within the target genome including deletion, insertion and translocations. Based on Codex Alimentarius claims, these induced genetic alterations can also be detected in normal plants breeding rather than GM foods. Among various Food and Drug organizations, Food Standards Australia New Zealand (FSANZ) has restricted using of GM foods following further assessments on human and animal studies⁶⁸.

Taken together, GM foods had great effects on human health and agriculture industry improvements. Despite of some controversial reports on possible side effects of GM foods in animal studies and on biodiversity of plants, review of the present studies is indicating that these impacts are mostly transient and like other genetic changes would be lost or existed by natural selection. However, further large scale and long-term analysis on both animal and human are warranted to define clearly the status of GM foods in nature food chains.

References

- 1 Reed, A. J. *et al.* Delayed Ripening Tomato Plants Expressing the Enzyme 1-Aminocyclopropane-1-carboxylic Acid Deaminase. 1. Molecular Characterization, Enzyme Expression, and Fruit Ripening Traits.



- Journal of Agricultural and Food Chemistry* 43, 1954-1962, doi:10.1021/jf00055a036 (1995).
- 2 Jackson, D. A., Symons, R. H. & Berg, P. Biochemical method for inserting new genetic information into DNA of Simian Virus 40: circular SV40 DNA molecules containing lambda phage genes and the galactose operon of *Escherichia coli*. *Proceedings of the National Academy of Sciences of the United States of America* 69, 2904-2909 (1972).
 - 3 Christou, P. & Twyman, R. M. The potential of genetically enhanced plants to address food insecurity. *Nutrition research reviews* 17, 23-42, doi:10.1079/nrr200373 (2004).
 - 4 Byrnes, B. H. & Bumb, B. L. Population Growth, Food Production and Nutrient Requirements. *Journal of Crop Production* 1, 1-27, doi:10.1300/J144v01n02_01 (1998).
 - 5 Cho, S. W., Lee, S. & Shin, W. The X-ray structure of *Aspergillus aculeatus* polygalacturonase and a modeled structure of the polygalacturonase-octagalacturonate complex. *Journal of molecular biology* 311, 863-878, doi:10.1006/jmbi.2001.4919 (2001).
 - 6 Murtaza, M. A., Ur-Rehman, S., Anjum, F. M., Huma, N. & Hafiz, I. Cheddar cheese ripening and flavor characterization: a review. *Critical reviews in food science and nutrition* 54, 1309-1321, doi:10.1080/10408398.2011.634531 (2014).
 - 7 Hafsa, A. B., Nabi, N., Zellama, M. S., Said, K. & Chaouachi, M. A new specific reference gene based on growth hormone gene (GH1) used for detection and relative quantification of Aquadvantage(R) GM salmon (*Salmo salar* L.) in food products. *Food chemistry* 190, 1040-1045, doi:10.1016/j.foodchem.2015.06.064 (2016).
 - 8 Gibbon, B. C. & Larkins, B. A. Molecular genetic approaches to developing quality protein maize. *Trends in genetics : TIG* 21, 227-233, doi:10.1016/j.tig.2005.02.009 (2005).
 - 9 Lyons, G. H., Stangoulis, J. C. & Graham, R. D. Exploiting micronutrient interaction to optimize biofortification programs: the case for inclusion of selenium and iodine in the HarvestPlus program. *Nutrition reviews* 62, 247-252 (2004).
 - 10 White, P. J. & Broadley, M. R. Biofortifying crops with essential mineral elements. *Trends in plant science* 10, 586-593, doi:10.1016/j.tplants.2005.10.001 (2005).
 - 11 Dai, J. L., Zhu, Y. G., Zhang, M. & Huang, Y. Z. Selecting iodine-enriched vegetables and the residual effect of iodate application to soil. *Biological trace element research* 101, 265-276, doi:10.1385/bter:101:3:265 (2004).
 - 12 Hartikainen, H. Biogeochemistry of selenium and its impact on food chain quality and human health. *Journal of trace elements in medicine and biology : organ of the Society for Minerals and Trace Elements (GMS)* 18, 309-318, doi:10.1016/j.jtemb.2005.02.009 (2005).
 - 13 Frossard, E., Bucher, M., Mächler, F., Mozafar, A. & Hurrell, R. Potential for increasing the content and bioavailability of Fe, Zn and Ca in plants for human nutrition. *Journal of the Science of Food and Agriculture* 80, 861-879, doi:10.1002/(SICI)1097-0010(20000515)80:7<861::AID-JSFA601>3.0.CO;2-P (2000).
 - 14 Bouis, H. E. Micronutrient fortification of plants through plant breeding: can it improve nutrition in man at low cost? *The Proceedings of the Nutrition Society* 62, 403-411 (2003).
 - 15 Bouis, H. E., Hotz, C., McClafferty, B., Meenakshi, J. V. & Pfeiffer, W. H. Biofortification: a new tool to reduce micronutrient malnutrition. *Food and nutrition bulletin* 32, S31-40, doi:10.1177/15648265110321s105 (2011).
 - 16 Gregorio, G. B. Progress in breeding for trace minerals in staple crops. *The Journal of nutrition* 132, 500S-502S (2002).
 - 17 Sainz, M. *et al.* Leghemoglobin is nitrated in functional legume nodules in a tyrosine residue within the heme cavity by a nitrite/peroxide-dependent mechanism. *The Plant journal : for cell and molecular biology* 81, 723-735, doi:10.1111/tpj.12762 (2015).
 - 18 Navascues, J. *et al.* Leghemoglobin green derivatives with nitrated hemes evidence production of highly reactive nitrogen species during aging of legume nodules. *Proceedings of the National Academy of Sciences of the United States of America* 109, 2660-2665, doi:10.1073/pnas.1116559109 (2012).
 - 19 Cong, L. *et al.* Multiplex genome engineering using CRISPR/Cas systems. *Science (New York, N.Y.)* 339, 819-823, doi:10.1126/science.1231143 (2013).
 - 20 Barampuram, S. & Zhang, Z. J. Recent advances in plant transformation. *Methods in molecular biology (Clifton, N.J.)* 701, 1-35, doi:10.1007/978-1-61737-957-4_1 (2011).
 - 21 DeMayo, F. J. & Spencer, T. E. CRISPR bacon: a sizzling technique to generate genetically engineered pigs. *Biology of reproduction* 91, 79, doi:10.1095/biolreprod.114.123935 (2014).
 - 22 Schultz, J. C., Appel, H. M., Ferrieri, A. P. & Arnold, T. M. Flexible resource allocation

- during plant defense responses. *Frontiers in plant science* 4, 324, doi:10.3389/fpls.2013.00324 (2013).
- 23 Bikker, P., Jongbloed, A. W. & Thissen, J. T. Meta-analysis of effects of microbial phytase on digestibility and bioavailability of copper and zinc in growing pigs. *Journal of animal science* 90 Suppl 4, 134-136, doi:10.2527/jas.53798 (2012).
 - 24 Al-Babili, S., Hoa, T. T. & Schaub, P. Exploring the potential of the bacterial carotene desaturase CrtI to increase the beta-carotene content in Golden Rice. *Journal of experimental botany* 57, 1007-1014, doi:10.1093/jxb/erj086 (2006).
 - 25 Hsieh, H. M., Liu, W. K. & Huang, P. C. A novel stress-inducible metallothionein-like gene from rice. *Plant molecular biology* 28, 381-389 (1995).
 - 26 Zhang, J. *et al.* Metallothionein 2 (SaMT2) from *Sedum alfredii* Hance confers increased Cd tolerance and accumulation in yeast and tobacco. *PloS one* 9, e102750, doi:10.1371/journal.pone.0102750 (2014).
 - 27 Frey, J. Biological safety concepts of genetically modified live bacterial vaccines. *Vaccine* 25, 5598-5605, doi:10.1016/j.vaccine.2006.11.058 (2007).
 - 28 Clarke, J. L., Waheed, M. T., Lossl, A. G., Martinussen, I. & Daniell, H. How can plant genetic engineering contribute to cost-effective fish vaccine development for promoting sustainable aquaculture? *Plant molecular biology* 83, 33-40, doi:10.1007/s11103-013-0081-9 (2013).
 - 29 Pelosi, A. *et al.* The release and induced immune responses of a plant-made and delivered antigen in the mouse gut. *Current drug delivery* 8, 612-621 (2011).
 - 30 Reboldi, A. *et al.* IgA production requires B cell interaction with subepithelial dendritic cells in Peyer's patches. *Science (New York, N.Y.)* 352, aaf4822, doi:10.1126/science.aaf4822 (2016).
 - 31 Concha, C. *et al.* Disease Prevention: An Opportunity to Expand Edible Plant-Based Vaccines? *Vaccines* 5, doi:10.3390/vaccines5020014 (2017).
 - 32 Lamichhane, A., Azegamia, T. & Kiyono, H. The mucosal immune system for vaccine development. *Vaccine* 32, 6711-6723, doi:10.1016/j.vaccine.2014.08.089 (2014).
 - 33 Clarke, J. L. *et al.* Lettuce-produced hepatitis C virus E1E2 heterodimer triggers immune responses in mice and antibody production after oral vaccination. *Plant biotechnology journal*, doi:10.1111/pbi.12743 (2017).
 - 34 Gregory, J. A. & Mayfield, S. P. Developing inexpensive malaria vaccines from plants and algae. *Applied microbiology and biotechnology* 98, 1983-1990, doi:10.1007/s00253-013-5477-6 (2014).
 - 35 Tacket, C. O. Plant-based vaccines against diarrheal diseases. *Transactions of the American Clinical and Climatological Association* 118, 79-87 (2007).
 - 36 Tacket, C. O. Plant-based oral vaccines: results of human trials. *Current topics in microbiology and immunology* 332, 103-117, doi:10.1007/978-3-540-70868-1_6 (2009).
 - 37 Mason, H. S. *et al.* Expression of Norwalk virus capsid protein in transgenic tobacco and potato and its oral immunogenicity in mice. *Proceedings of the National Academy of Sciences of the United States of America* 93, 5335-5340 (1996).
 - 38 Herbst-Kralovetz, M., Mason, H. S. & Chen, Q. Norwalk virus-like particles as vaccines. *Expert review of vaccines* 9, 299-307, doi:10.1586/erv.09.163 (2010).
 - 39 Joung, Y. H. *et al.* The Last Ten Years of Advancements in Plant-Derived Recombinant Vaccines against Hepatitis B. *International journal of molecular sciences* 17, doi:10.3390/ijms17101715 (2016).
 - 40 Santi, L., Huang, Z. & Mason, H. Virus-like particles production in green plants. *Methods (San Diego, Calif.)* 40, 66-76, doi:10.1016/j.ymeth.2006.05.020 (2006).
 - 41 Rademacher, T. *et al.* Recombinant antibody 2G12 produced in maize endosperm efficiently neutralizes HIV-1 and contains predominantly single-GlcNAc N-glycans. *Plant biotechnology journal* 6, 189-201, doi:10.1111/j.1467-7652.2007.00306.x (2008).
 - 42 Rubio-Infante, N. *et al.* A Plant-Derived Multi-HIV Antigen Induces Broad Immune Responses in Orally Immunized Mice. *Molecular biotechnology* 57, 662-674, doi:10.1007/s12033-015-9856-3 (2015).
 - 43 Biemelt, S., Sonnewald, U., Galmbacher, P., Willmitzer, L. & Muller, M. Production of human papillomavirus type 16 virus-like particles in transgenic plants. *Journal of virology* 77, 9211-9220 (2003).
 - 44 Greco, R. *et al.* Production of recombinant HIV-1/HBV virus-like particles in *Nicotiana tabacum* and *Arabidopsis thaliana* plants for a bivalent plant-based vaccine. *Vaccine* 25, 8228-8240, doi:10.1016/j.vaccine.2007.09.061 (2007).
 - 45 Phoolcharoen, W. *et al.* A nonreplicating subunit vaccine protects mice against lethal



- Ebola virus challenge. *Proceedings of the National Academy of Sciences of the United States of America* 108, 20695-20700, doi:10.1073/pnas.1117715108 (2011).
- 46 Nemchinov, L. G. *et al.* Development of a plant-derived subunit vaccine candidate against hepatitis C virus. *Archives of virology* 145, 2557-2573 (2000).
- 47 Aboul-Ata, A. A. *et al.* Plant-based vaccines: novel and low-cost possible route for Mediterranean innovative vaccination strategies. *Advances in virus research* 89, 1-37, doi:10.1016/b978-0-12-800172-1.00001-x (2014).
- 48 Chichester, J. A. *et al.* A plant-produced protective antigen vaccine confers protection in rabbits against a lethal aerosolized challenge with *Bacillus anthracis* Ames spores. *Human vaccines & immunotherapeutics* 9, 544-552 (2013).
- 49 Rosales-Mendoza, S. & Tello-Olea, M. A. Carrot cells: a pioneering platform for biopharmaceuticals production. *Molecular biotechnology* 57, 219-232, doi:10.1007/s12033-014-9837-y (2015).
- 50 Yao, J., Weng, Y., Dickey, A. & Wang, K. Y. Plants as Factories for Human Pharmaceuticals: Applications and Challenges. *International journal of molecular sciences* 16, 28549-28565, doi:10.3390/ijms161226122 (2015).
- 51 Kytidou, K. *et al.* Human Alpha Galactosidases Transiently Produced in *Nicotiana benthamiana* Leaves: New Insights in Substrate Specificities with Relevance for Fabry Disease. *Frontiers in plant science* 8, 1026, doi:10.3389/fpls.2017.01026 (2017).
- 52 Jobsri, J. *et al.* Plant virus particles carrying tumour antigen activate TLR7 and Induce high levels of protective antibody. *PloS one* 10, e0118096, doi:10.1371/journal.pone.0118096 (2015).
- 53 Hefferon, K. Reconceptualizing cancer immunotherapy based on plant production systems. *Future science OA* 3, FSO217, doi:10.4155/fsoa-2017-0018 (2017).
- 54 Hilder, V. A., Gatehouse, A. M. R., Sheerman, S. E., Barker, R. F. & Boulter, D. A novel mechanism of insect resistance engineered into tobacco. *Nature* 330, 160-163 (1987).
- 55 Green, J. M. & Owen, M. D. Herbicide-resistant crops: utilities and limitations for herbicide-resistant weed management. *J Agric Food Chem* 59, 5819-5829, doi:10.1021/jf101286h (2011).
- 56 Iquebal, M. A. *et al.* Discovery of Putative Herbicide Resistance Genes and Its Regulatory Network in Chickpea Using Transcriptome Sequencing. *Frontiers in plant science* 8, 958, doi:10.3389/fpls.2017.00958 (2017).
- 57 Shoba, D. *et al.* Development and Genetic Characterization of A Novel Herbicide (Imazethapyr) Tolerant Mutant in Rice (*Oryza sativa* L.). *Rice (New York, N.Y.)* 10, 10, doi:10.1186/s12284-017-0151-8 (2017).
- 58 Yu, X. D. *et al.* RNAi-mediated plant protection against aphids. *Pest management science* 72, 1090-1098, doi:10.1002/ps.4258 (2016).
- 59 Saumet, A. & Lecellier, C. H. Anti-viral RNA silencing: do we look like plants? *Retrovirology* 3, 3, doi:10.1186/1742-4690-3-3 (2006).
- 60 Yang, H. *et al.* Review: biosafety assessment of Bt rice and other Bt crops using spiders as example for non-target arthropods in China. *Plant cell reports* 36, 505-517, doi:10.1007/s00299-017-2108-1 (2017).
- 61 Baktavachalam, G. B. *et al.* Transgenic maize event TC1507: Global status of food, feed, and environmental safety. *GM crops & food* 6, 80-102, doi:10.1080/21645698.2015.1054093 (2015).
- 62 Szenasi, A., Palinkas, Z., Zalai, M., Schmitz, O. J. & Balog, A. Short-term effects of different genetically modified maize varieties on arthropod food web properties: an experimental field assessment. *Scientific reports* 4, 5315, doi:10.1038/srep05315 (2014).
- 63 Cook-Patton, S. C., McArt, S. H., Parachnowitsch, A. L., Thaler, J. S. & Agrawal, A. A. A direct comparison of the consequences of plant genotypic and species diversity on communities and ecosystem function. *Ecology* 92, 915-923 (2011).
- 64 Romeis, J., Meissle, M. & Bigler, F. Transgenic crops expressing *Bacillus thuringiensis* toxins and biological control. *Nature biotechnology* 24, 63-71, doi:10.1038/nbt1180 (2006).
- 65 Warwick, S. I., Becki.e., H. J. & Hall, L. M. Gene flow, invasiveness, and ecological impact of genetically modified crops. *Annals of the New York Academy of Sciences* 1168, 72-99, doi:10.1111/j.1749-6632.2009.04576.x (2009).
- 66 Chapman, M. A. & Burke, J. M. Letting the gene out of the bottle: the population genetics of genetically modified crops. *The New phytologist* 170, 429-443, doi:10.1111/j.1469-8137.2006.01710.x (2006).
- 67 Liu, N. *et al.* Effect on soil chemistry of genetically modified (GM) vs. non-GM maize.



- GM crops* 1, 157-161, doi:10.4161/gmcr.1.3.12810 (2010).
- 68 Safety and nutritional assessment of GM plants and derived food and feed: the role of animal feeding trials. *Food and chemical toxicology : an international journal published for the British Industrial Biological Research Association* 46 Suppl 1, S2-70, doi:10.1016/j.fct.2008.02.008 (2008).