

## A narrative review on the application of sanitizers in the COVID-19 pandemic

Vahid Ranaei <sup>1</sup>, Leyla Alizadeh <sup>2</sup>, Zahra Pilevar <sup>3</sup>, Teamur Aghamolaei <sup>4\*</sup>, Hedayat Hosseini <sup>5\*</sup>

<sup>1</sup> Student Research Committee, Faculty of Health, Hormozgan University of Medical Sciences, Bandar Abbas, Iran

<sup>2</sup> Department of Food Quality Control and Hygiene, Science and Research Branch, Islamic Azad University, Tehran, Iran

<sup>3</sup> School of Health, Arak University of Medical Sciences, Arak, Iran

<sup>4</sup> Cardiovascular Research Center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran

<sup>5</sup> Department of Food Science and Technology, National Nutrition and Food Technology Research Institute, Faculty of Nutrition Sciences and Food Technology, Shahid Beheshti University of Medical Sciences, Tehran, Iran

### ARTICLE INFO

#### Review Article

#### Article history:

Received 22 December 2021

Revised 19 February 2022

Accepted 27 February 2022

Available online 20 March 2022

#### Keywords:

COVID-19 pandemic

Disinfectant

Sanitizers

Hand hygiene

### ABSTRACT

At the beginning of 2020, a new coronavirus, namely SARS-CoV-2, has started to spread worldwide leading to the so-called COVID-19 pandemic. The present study aims to review the effectiveness of disinfectants in the prevention and control of the COVID-19 pandemic. In this article, after an introduction to the COVID-19 pandemic, the role of disinfectants in the prevention and control of the COVID-19 pandemic is discussed. To identify the articles, English databases including Google Scholar, Scopus, Science Direct, Web of Science, and PubMed from December 2020 to November 2020 were examined using keywords. There is also a manual search in journals, a collection of abstracts in conferences, and conferences and dissertations. Hand hygiene is a well-accepted principle in preventing the transmission of most infectious diseases. Given that many formulations of sanitizers can be effective against COVID-19, it may alter the integrity and function of the skin barrier and increase the risk of dermatitis on the hands.

© 2022, Science and Research Branch, Islamic Azad University. All rights reserved.

### 1. Introduction

World Health Organization (WHO) declared COVID-19 as a pandemic on March 11, 2020. The global infection rate of COVID-19 is increasing sharply. Coronaviruses (CoV) represent a diverse and large family of enveloped and single-stranded RNA viruses (1). The structure of this family can cause a variety of illnesses in humans and animals including Severe Acute Respiratory Syndrome (SARS), Middle East respiratory syndrome (MERS), and COVID-19 (2). Coronaviruses comprise alpha, gamma, beta, and delta groups with 22 subclasses and 40 species (3). The structure of COVID-19 is shown in Fig. 1. The spike proteins are responsible for the synthesis of RNA, whereas membrane and envelope proteins are associated with viral assembly (4). The common symptoms of COVID-19 disease are pneumonia, acute respiratory disease, fever, and dry cough. However, this disease may infect people in an asymptomatic form. Given that millions of people are receiving the vaccine, however, there

are no medications to treat COVID-19 disease and a new highly contagious mutated form of coronavirus is emerging. Several mitigation policies including city lockdowns and social distancing have been implemented by the governments (4). Personal and social preventive measures slow down the super spread of the pandemic until effective drugs become available and vaccination is done (5). WHO has recommended people have a healthy lifestyle, strengthen their immune system, wear masks and disinfect hands with soap or alcohol-based sanitizers. In this review, the role and mechanism of disinfectants against coronavirus are briefly discussed.

### 2. Use of hand hygiene products

Hand hygiene as a principle can decrease the spread of respiratory diseases and the COVID-19 pandemic (6, 7). Given that hand hygiene practices are easy and inexpensive, however, compliance with handwashing differs from approximately 2 to 78% among various populations and

Corresponding authors: \*Cardiovascular research Center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran.  
E-mail address: [teaghamolaei@gmail.com](mailto:teaghamolaei@gmail.com) (Teamur Aghamolaei).

\*Department of Food Science and Technology, National Nutrition and Food Technology Research Institute, Faculty of Nutrition Sciences and Food Technology, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

E-mail address: [hedayat@sbmu.ac.ir](mailto:hedayat@sbmu.ac.ir) (Hedayat Hosseini).

contexts (8, 9). Perception and effectiveness of hand hygiene performance are associated with many factors including gender distinction (10, 11). Girls have shown 1.12 times the likelihood of excellent handwashing compared to boys (12).




Fig. 1. Structure of COVID-19 (4).

Soap as a fatty acid salt has cleansing properties that remove viruses by disrupting intracellular and membrane lipids. However, frequent washing with water and soap can lead to irritation and skin sensitivity due to damage to keratin proteins and removal of intracellular lipids which are in the stratum corneum of the skin (13, 14). However, several factors other than harsh soaps and potent detergents such as water hardness, rubbing with rough paper towels, wet work, low humidity, working with irritants, and hot or very cold water can affect the function of the skin barrier (15, 16). American Contact Dermatitis Society declared that irritant contact dermatitis and allergic contact dermatitis increases during the COVID-19 pandemic by stringent hand hygiene (14). As reported, health care workers wash their hands more than ten times every day and only a few of them use moisturizers after washing their hands which leads to hand dermatitis (6, 17, 18). As reported by Derksen et al. (19) hand hygiene performance has increased from 47 to 95 percent after lockdown during the COVID-19 pandemic. Taghrir et al. (20) study assessed the knowledge, self-report preventive behaviors (hand washing and surface disinfection), and risk perception related to COVID-19, among Iranian medical students in the first week after the Covid-19 outbreak in Iran. This cross-sectional study was conducted from February 26 to 28, 2020. Participants were Iranian medical students (fifth to the seventh year) whose knowledge, preventive behaviors (hand washing and surface disinfection), and their perception of COVID-19 risk were assessed using an online questionnaire. The questionnaire consisted of 26 questions including 15 questions about knowledge and awareness related to COVID-19, 9 questions about preventive measures (hand washing and disinfection of surfaces), and 2 questions about risk perception related to COVID-19. The validity and reliability of the questionnaire were satisfactory. A total of 240 medical students completed the questionnaire. The results showed that the mean age of the participants was 23 years. The mean of correct knowledge responses was 96.86% and 79.60% of the participants had high awareness. The average rate of preventive behaviors was

94.47%. and 94.2% had good performance in preventive behaviors (hand washing and surface disinfection). The cumulative risk perception was 4.08 out of 8, which was moderate. There was a statistically significant difference in risk perception between interns and trainees as well as between those trained in the emergency and non-emergency departments. There was a negative correlation between preventive behaviors (hand washing and surface disinfection) and risk perception. There was also a high level of knowledge and awareness related to COVID-19 and self-reported preventive behaviors (hand washing and surface disinfection) and risk perception among Iranian medical students. The study by Moore et al. (21) aimed to investigate the effect of the COVID-19 pandemic on hand health performance (HHP) in acute care hospitals in the United States. The increase is discussed and then decreases as it progresses. In this paper, hand hygiene performance (HHP) rates were assessed using an automatic hand hygiene monitoring system in 74 adult admission wards in 7 hospitals and 10 pediatric admission units in 2 pediatric hospitals. A regression model was used to estimate hand hygiene performance rates at 10 weeks leading up to a significant COVID-19 event (e.g., school closure) and for 10 weeks thereafter. Various analyzes were performed and several scenarios were studied to elucidate the effect of the parameters on the possible outcomes. Data from an automated hand hygiene monitoring system showed that hand hygiene performance in the early stages of the epidemic has changed in several directions. Three effects emerged that were all significant at  $p < 0.01$ . The results showed that the average rate of average hand hygiene performance in the months before the school closure related to the start of the epidemic increased from 46 to 56. At the time of school closure, there was a 6% upward shift. The hand hygiene performance rate remained more than 60% for 4 weeks before decreasing to 54% at the end of the study period. The study conducted by Derxen et al. (19) studied the adherence and evaluating the determining factors in obstetric and gynecological hospitals during and after COVID-19 in Germany. In this article, a total of 267 behavioral observations were made in two hospitals of the German University of Obstetrics and Gynecology in three time periods (before the COVID-19 pandemic, increased awareness and careful precautions). In addition, 115 health workers answered questionnaires about the social cognitive determinants of hand health behavior. Multiple regression was used to analyze the relationships. Findings showed that adherence to hand hygiene recommendations increased from 47% of cases before the COVID-19 pandemic to 95% just before the restrictions (although simple anti-epidemic measures were taken). Self-efficacy was associated with the intention to follow hand disinfection instructions ( $b = 0.397$ ,  $p < 0.001$ ). Obstetricians and gynecologists appear to adapt their hand disinfection behavior to prevent infections in the COVID-19 pandemic. To further improve preventive interventions, cognitive-social determinants, especially behavioral intent and self-efficacy, must be considered. The study by Chen et al. (12) aimed to understand the health status of hand cuffs and masks among primary school students during

the 2019 Coronavirus (COVID-19) pandemic in Wuhan, China. In this paper, logistic regression analysis was used to identify risk factors affecting handwashing behaviors and mask use. The results showed that 42.05% of primary school students showed good handwashing behavior, while 51.60% had good behavior in using the mask. Gender, educational background, educational background, father's job, mother's educational background and time to complete the survey were significantly related to hand health, while educational background, mother's education was related to mask-wearing behavior. Hand-washing and masking behaviors among primary school students are influenced by gender, grade, and other factors, so parents should follow their instructions, while governments should provide general advice on hand-washing and masking behaviors. The synthetic detergents which are a mixture of petrolatum and surfactants have a similar pH to the skin (5.5-7) and contain less than 10 percent of soap (13). Synthetic detergents such as glucosides, cocamide diethanolamine, and sodium cocoyl glutamate can disrupt the viral membrane of lipid enveloped viruses (22, 23). The concentration of surfactants is an important factor for irritation, inflammation, and skin xerosis (13, 14). Antiseptic handwashes are soaps and synthetic detergents that have antimicrobial agents that disrupt the integrity of the viral membrane. For example, alcohol denatures proteins or povidone iodine impairs protein synthesis by penetrating to cells and inactivating the cellular replication (14). The hand wipes contain antimicrobial agents such as alcohol and benzethonium chloride which disrupt the integrity and viral membrane of enveloped viruses (14). It worth mentioning that surface wipes should not be used for skin wipes as they contain harsh antimicrobial agents like N-alkyl dimethyl benzyl ammonium chloride (24). The Centers for Disease Control and Prevention has recommended using alcohol-based sanitizers that contain moisturizers and emollients with low allergenicity instead of harsh soaps and synthetic detergents to prevent contact dermatitis (25, 26). The alcohol-based sanitizers contain ethanol, isopropanol, and hydrogen peroxide portfolio (27). The high-touch surfaces should be disinfected. After cleaning and removal of organic matters from surfaces by brushing and using soap and water, disinfectants can be used to kill any residual microorganisms (28, 29). It should be noted that fogging/spraying of chemical disinfectants such as quaternary ammonium compounds, chlorine-based agents, and formaldehyde for removal of COVID-19 from environmental surfaces is not recommended due to their adverse health effects (30). As reported by Slaughter et al. (31), alcohol-based sanitizers can also have toxic adverse impacts on the environment after evaporation. In the first five months of 2020, the Pesticide Control Association reported 9504 cases of contact with alcoholic disinfectants in children under 12 years of age. Even small amounts of alcohol can cause alcohol poisoning in children, causing dizziness, vomiting, drowsiness, and in severe cases, respiratory arrest and death (32). Disinfectants including isopropanol, hydrogen peroxide, formaldehyde, iodophors, and ethanol can cause respiratory and non-respiratory problems such as gastrointestinal effects,

skin and eye irritation, dermatitis, liver damage, and central nervous system depression (33, 34). For example, skin absorption of ethanol-based disinfectants via occupational exposure and frequent use of hand sanitizers can cause poisoning (34). On the other hand, skin as a barrier to viruses and bacteria can become more susceptible to many infections and diseases due to the use of disinfectants that remove skin oil and increase skin permeability (28). It has been reported that excessive use of alcohol-based disinfectants can increase the risk of viral outbreaks (35). To avoid these adverse health problems, technologies without direct exposure to disinfectants such as irradiation with UV, aerosolized hydrogen peroxide, gaseous ozone, and pulsed xenon have been proposed (36). It is recommended that if cleaners are in direct contact with disinfectants and contaminated surfaces, personal protective equipment (PPE) like gowns, masks, and face shields should be used (30). As COVID-19 can be transmitted via air, it is important to use PPE for safety. The type and concentration of disinfectant for high-touch surfaces like food preparation areas should be carefully selected to prevent adverse health effects and avoid damaging the food surfaces (28).

### **3. Mechanism of action of some disinfectants on coronavirus and other enveloped viruses**

The spherical coronavirus as an enveloped virus with an outer lipid layer is more sensitive to disinfectants than non-enveloped viruses (28). The antigenic proteins of the outer surface envelope attach to specific receptors of the host. The virus enters the host cell and replicates by integrating with the genetic RNA code (37). The alcohol-based disinfectants containing ethanol penetrate the membrane and denatures the protein tertiary structure by the breakdown of hydrogen bonds (38). Ethanol is more effective against hydrophilic viruses such as COVID-19, rotavirus, and human immunodeficiency virus (HIV) than isopropyl alcohol (28, 39). Isopropyl alcohol and ethanol exhibit 70-90% biocide properties against COVID-19 (30 sec) (38). Alcohol-based hand sanitizers consist of 60-95% alcohol (40). The carcinogenicity of alcohol-based disinfectants through skin absorption is still unknown (41). Alcohol toxicity can cause hypophosphatemia, hypomagnesemia, hypocalcemia, myoglobinuria, respiratory depression, and acute liver damage (42, 43). Determination of the toxic concentration of ethanol-based hand rub is difficult due to different tolerance levels and reactions to ethanol in different people (32). It should be noted that oral consumption of alcohol through alcohol-based sanitizers, alcohol-containing cosmetics, and mouthwash can cause toxicity (44). Dermal exposure to ethanol leads to allergic responses and drying and cracking of the skin which can be reduced by the application of combined components with ethanol such as panthenol to regenerate the skin barrier or glycerol for the hydration of skin (45). Peroxide-based disinfectants such as peroxy-acetic acid and hydrogen peroxide affect nucleic acid, protein, and lipid membrane of viruses via producing hydroxyl free radicals (46). Hydrogen peroxide at 1-3 % levels oxidizes

disulfide bonds of thiol groups in proteins and inactivates SARS-CoV (28). The hypochlorous acid (HOCl) as a solution of sodium and calcium hypochlorite exhibits extensive antiviral activity. Given that hypochlorite reacts and inactivates in exposure to organic matters, surfaces should be mechanically cleaned with detergents or soap and water before application of these disinfectants (47). The activity of chlorine solution is higher at lower pH values. However, the shelf life of low pH chlorine solution is shorter than high pH chlorine solution and should be prepared fresh for use (48). WHO has declared that chlorine-based disinfectants kill 99% of viruses (49). Alkyl dimethyl ammonium chloride biocide as a quaternary ammonium compound in a low concentration of 1 % is efficient against COVID-19 in less than 1 minute (38). As reported by Kariwa et al. (50) formaldehyde and glutaraldehyde can alkylate the bacterial proteins and nucleic acid at concentrations of 0.5-3%. Povidone-iodine as an antiseptic agent for viruses can damage nucleic acid and break down the disulfide bonds in proteins which can inactivate SARS-CoV at a concentration of less than 1% (51). Health care workers have improved their hand hygiene behaviors to protect themselves and their families from this COVID-19 pandemic (52). Given that emphasizing the importance of hand hygiene improves hand hygiene performance (21), in the food industry, perception of personal risk and self-protection is not a driver for hand hygiene and there should be other reasons for a good hand hygiene performance (53).

#### 4. Conclusion

It is recommended to wash hands with antimicrobial soaps to eliminate possible infection with COVID-19 disease. However, the excessive use of hand sanitizers as a preventive measure against COVID-19 is not environmentally friendly and is dangerous to human health. To date, no drugs are available to combat COVID 19 infection, and the rate of infection is rising sharply worldwide. The World Health Organization (WHO) has proposed only vaccination and preventative measures and a healthy lifestyle with an effective immune system to combat and stay safe from the pandemic. The World Health Organization recommends alcohol-based hand sanitizers for recurrent hand hygiene, which are mainly made from ethanol, isopropyl alcohols, and hydrogen peroxides with various compounds. The compounds of these substances may be toxic to human health and the environment, and if used improperly, these chemicals have toxic and dangerous effects on the environment that are released during evaporation. In addition, repeated use of hand sanitizers has increased the risk of antimicrobial resistance and other viral diseases.

#### References

- Payne S. Family coronaviridae. *Viruses*. 2017;149.
- Park SE. Epidemiology, virology, and clinical features of severe acute respiratory syndrome-coronavirus-2 (SARS-CoV-2; Coronavirus Disease-19). *Clinical and Experimental Pediatrics*. 2020;63(4):119.
- Ranaei V, Pilevar Z, Hosseini H. Food safety practices in COVID-19 pandemic. *Journal of Food Quality and Hazards Control*. 2020;7(3):116-8.
- Ranaei V, Pilevar Z, Neyestani TR. Can Raising Vitamin D Status Slow Down Covid-19 Waves? *Nutrition and Food Sciences Research*. 2021;8(1):1-3.
- Ranaei V, Yarmohammadi S, Alizadeh L, Dadipoor S, Pilevar Z, Aghamolaei T. Emerging 2019-ncov disease: A narrative review on the personal and social preventive behaviors. *Food & Health*. 2021; 4(3):13-8.
- Kantor J. Behavioral considerations and impact on personal protective equipment use: Early lessons from the coronavirus (COVID-19) pandemic. *Journal of the American Academy of Dermatology*. 2020;82(5):1087-8.
- Fung ICH, Cairncross S. Effectiveness of handwashing in preventing SARS: a review. *Tropical Medicine & International Health*. 2006;11(11):1749-58.
- Hirai M, Graham JP, Mattson KD, Kelsey A, Mukherji S, Cronin AA. Exploring determinants of handwashing with soap in Indonesia: a quantitative analysis. *International Journal of Environmental Research and Public Health*. 2016;13(9):868.
- Ali MM, Verrill L, Zhang Y. Self-reported hand washing behaviors and foodborne illness: a propensity score matching approach. *Journal of Food Protection*. 2014;77(3):352-8.
- Rubin GJ, Amlôt R, Page L, Wessely S. Public perceptions, anxiety, and behaviour change in relation to the swine flu outbreak: cross sectional telephone survey. *BMJ*. 2009;339.
- Park J-H, Cheong H-K, Son D-Y, Kim S-U, Ha C-M. Perceptions and behaviors related to hand hygiene for the prevention of H1N1 influenza transmission among Korean university students during the peak pandemic period. *BMC Infectious Diseases*. 2010;10(1):1-8.
- Chen X, Ran L, Liu Q, Hu Q, Du X, Tan X. Hand hygiene, mask-wearing behaviors and its associated factors during the COVID-19 epidemic: A cross-sectional study among primary school students in Wuhan, China. *International Journal of Environmental Research and Public Health*. 2020;17(8):2893.
- Levin J, Miller R. A guide to the ingredients and potential benefits of over-the-counter cleansers and moisturizers for rosacea patients. *The Journal of Clinical and Aesthetic Dermatology*. 2011;4(8):31.
- Rundle CW, Presley CL, Militello M, Barber C, Powell DL, Jacob SE, et al. Hand hygiene during COVID-19: recommendations from the American Contact Dermatitis Society. *Journal of the American Academy of Dermatology*. 2020;83(6):1730-7.
- Abtahi-Naeini B. Frequent handwashing amidst the COVID-19 outbreak: prevention of hand irritant contact dermatitis and other considerations. *Health Science Reports*. 2020;3(2):e16.
- Danby SG, Brown K, Wigley AM, Chittock J, Pyae PK, Flohr C, et al. The effect of water hardness on surfactant deposition after washing and subsequent skin irritation in atopic dermatitis patients and healthy control subjects. *Journal of Investigative Dermatology*. 2018;138(1):68-77.
- Masood S, Tabassum S, Naveed S, Jalil P. COVID-19 Pandemic & skin care guidelines for health care professionals. *Pakistan Journal of Medical Sciences*. 2020;36(COVID19-S4):S115.
- Lan J, Song Z, Miao X, Li H, Li Y, Dong L, et al. Skin damage among health care workers managing coronavirus disease-2019. *Journal of the American Academy of Dermatology*. 2020;82(5):1215-6.
- Derksen C, Keller FM, Lippke S. Obstetric Healthcare Workers' Adherence to Hand Hygiene Recommendations during the COVID-19 Pandemic: Observations and Social-Cognitive Determinants. *Applied Psychology: Health and Well-Being*. 2020;12(4):1286-305.
- Taghbir MH, Borazjani R, Shiraly R. COVID-19 and Iranian medical students: a survey on their related-knowledge, preventive behaviors and risk perception. *Archives of Iranian Medicine*. 2020;23(4):249-54.
- Moore LD, Robbins G, Quinn J, Arbogast JW. The impact of COVID-19 pandemic on hand hygiene performance in hospitals. *American Journal of Infection Control*. 2021;49(1):30-3.
- Draeos ZD. The science behind skin care: Cleansers. *Journal of Cosmetic Dermatology*. 2018;17(1):8-14.
- Hellstern P, Solheim BG. The use of solvent/detergent treatment in pathogen reduction of plasma. *Transfusion Medicine and Hemotherapy*. 2011;38(1):65-70.

24. Mauleón C, Mauleón P, Chavarría E, De La Cueva P, Suárez R, Pablo L. Airborne contact dermatitis from n-alkyl dimethylbenzylammonium chloride and n-alkyl dimethylethyl-benzylammonium chloride in a detergent. *Contact Dermatitis*. 2006;55(5):311-2.
25. Xu S, Kwa M, Lohman ME, Evers-Meltzer R, Silverberg JI. Consumer preferences, product characteristics, and potentially allergenic ingredients in best-selling moisturizers. *JAMA Dermatology*. 2017;153(11):1099-105.
26. Swain SK. Practice of handwashing: An effective tool to control COVID-19 pandemic. *Journal of the Scientific Society*. 2021;48(2):60.
27. Kilpatrick C, Allegranzi B, Pittet D. WHO first global patient safety challenge: clean care is safer care, contributing to the training of health-care workers around the globe. *International Journal of Infection Control*. 2011;7(2).
28. Ghafoor D, Khan Z, Khan A, Ualiyeva D, Zaman N. Excessive use of disinfectants against COVID-19 posing potential threat to living beings. *Current Research in Toxicology*. 2021.
29. Chartier Y, Adams J, Bartram J. Essential environmental health standards in health care. Geneva: WHO. 2008.
30. Organization WH. Cleaning and disinfection of environmental surfaces in the context of COVID-19: interim guidance, 15 May 2020. World Health Organization; 2020.
31. Slaughter R, Mason R, Beasley D, Vale J, Schep L. Isopropanol poisoning. *Clinical Toxicology*. 2014;52(5):470-8.
32. Mahmood A, Eqan M, Pervez S, Alghamdi HA, Tabinda AB, Yasar A, et al. COVID-19 and frequent use of hand sanitizers; human health and environmental hazards by exposure pathways. *Science of the Total Environment*. 2020;742:140561.
33. Schyllert C, Rönmark E, Andersson M, Hedlund U, Lundbäck B, Hedman L, et al. Occupational exposure to chemicals drives the increased risk of asthma and rhinitis observed for exposure to vapours, gas, dust and fumes: a cross-sectional population-based study. *Occupational and Environmental Medicine*. 2016;73(10):663-9.
34. Salomone A, Bozzo A, Di Corcia D, Gerace E, Vincenti M. Occupational exposure to alcohol-based hand sanitizers: the diagnostic role of alcohol biomarkers in hair. *Journal of Analytical Toxicology*. 2018;42(3):157-62.
35. Vogel L. Hand sanitizers may increase norovirus risk. *Canadian Medical Association Journal*. 2011;183(12): E799–E800.
36. Weber DJ, Rutala WA, Anderson DJ, Chen LF, Sickbert-Bennett EE, Boyce JM. Effectiveness of ultraviolet devices and hydrogen peroxide systems for terminal room decontamination: focus on clinical trials. *American Journal of Infection Control*. 2016;44(5):e77-e84.
37. Kratzel A, Todt D, V'kovski P, Steiner S, Gultom M, Thao TTN, et al. Inactivation of severe acute respiratory syndrome coronavirus 2 by WHO-recommended hand rub formulations and alcohols. *Emerging Infectious Diseases*. 2020;26(7):1592.
38. Kampf G, Todt D, Pfaender S, Steinmann E. Persistence of coronaviruses on inanimate surfaces and their inactivation with biocidal agents. *Journal of Hospital Infection*. 2020;104(3):246-51.
39. Lantagne D, Wolfe M, Gallandat K, Opryszko M. Determining the Efficacy, safety and suitability of disinfectants to prevent emerging infectious disease transmission. *Water*. 2018;10(10):1397.
40. Barrett MJ, Babi FE. Alcohol-based hand sanitiser: a potentially fatal toy. *Medical Journal of Australia*. 2015;203(1):43-4.
41. Lachenmeier DW. Safety evaluation of topical applications of ethanol on the skin and inside the oral cavity. *Journal of Occupational Medicine and Toxicology*. 2008;3(1):1-16.
42. Wilson M, Guru P, Park J. Recurrent lactic acidosis secondary to hand sanitizer ingestion. *Indian Journal of Nephrology*. 2015;25(1):57.
43. Bouthoorn SH, Van der Ploeg T, Van Erkel NE, Van der Lely N. Alcohol intoxication among Dutch adolescents: acute medical complications in the years 2000-2010. *Clinical Pediatrics*. 2011;50(3):244-51.
44. Vonghia L, Leggio L, Ferrulli A, Bertini M, Gasbarrini G, Addolorato G, et al. Acute alcohol intoxication. *European Journal of Internal Medicine*. 2008;19(8):561-7.
45. Kampf G, Muscatiello M, Häntschel D, Rudolf M. Dermal tolerance and effect on skin hydration of a new ethanol-based hand gel. *Journal of Hospital Infection*. 2002;52(4):297-301.
46. Knotzer S, Kindermann J, Modrof J, Kreil TR. Measuring the effectiveness of gaseous virus disinfectants. *Biologicals*. 2015;43(6):519-23.
47. Aline K-SA, Adou-Assoumou M, Djoilé SX, Diemer F, Gurgel M. The effects of sodium hypochlorite on organic matters: Influences of concentration, renewal frequency and contact area. *Iranian Endodontic Journal*. 15(1):18-22.
48. Iqbal Q, Lubeck-Schricker M, Wells E, Wolfe MK, Lantagne D. Shelf-life of chlorine solutions recommended in Ebola virus disease response. *PLoS One*. 2016;11(5):e0156136.
49. Jayaprakash A, Srividhya Muralidharan PJ, Devendran G. Potent Action of Disinfectant in Targeting Spread of COVID-19 Using Drones. *International Journal of Research in Engineering, Science and Management*. 2020;3(5):808-10.
50. Kariwa H, Fujii N, Takashima I. Inactivation of SARS coronavirus by means of povidone-iodine, physical conditions and chemical reagents. *Dermatology*. 2006;212(Suppl. 1):119-23.
51. Eggers M, Koburger-Janssen T, Eickmann M, Zorn J. In vitro bactericidal and virucidal efficacy of povidone-iodine gargle/mouthwash against respiratory and oral tract pathogens. *Infectious Diseases and Therapy*. 2018;7(2):249-59.
52. Whitby M, McLaws M-L, Ross MW. Why healthcare workers don't wash their hands: a behavioral explanation. *Infection Control & Hospital Epidemiology*. 2006;27(5):484-92.
53. Erasmus V, Brouwer W, Van Beeck E, Oenema A, Daha T, Richardus JH, et al. A qualitative exploration of reasons for poor hand hygiene among hospital workers lack of positive role models and of convincing evidence that hand hygiene prevents cross-infection. *Infection Control & Hospital Epidemiology*. 2009;30(5):415-9.